

# Measures on perfect PAC fields with pro-cyclic Galois group

I. Halupczok

Ecole Normale Supérieure  
rue d'Ulm, Paris

Modnet Antalya Conference, 2006

# Introduction

- Emerged from work group in Lyon (France) with Thomas Blossier, Amador Martin Pizarro and Juan Pons Llopis.
- Answer a question of Hrushovski on “measures on fields with bounded Galois groups”.

# Outline

- 1 Measures on pseudo-finite fields
- 2 How to generalize?
- 3 Results

# Outline

- ① Measures on pseudo-finite fields
- ② How to generalize?
- ③ Results

# Outline

- ① Measures on pseudo-finite fields
- ② How to generalize?
- ③ Results

# Outline

- 1 Measures on pseudo-finite fields
- 2 How to generalize?
- 3 Results

# Situation for pseudo-finite fields

- Let  $K$  be a pseudo-finite field.
- Want to understand definable sets  $X \subset K^n$ .
- Method: Associate **invariants** to  $X$ :
  - Dimension  $\dim(X)$
  - Measure  $\mu(X)$

# Situation for pseudo-finite fields

- Let  $K$  be a pseudo-finite field.
- Want to understand definable sets  $X \subset K^n$ .
- Method: Associate **invariants** to  $X$ :
  - Dimension  $\dim(X)$
  - Measure  $\mu(X)$



# Situation for pseudo-finite fields

- Let  $K$  be a pseudo-finite field.
- Want to understand definable sets  $X \subset K^n$ .
- Method: Associate **invariants** to  $X$ :
  - Dimension  $\dim(X)$
  - Measure  $\mu(X)$

# Situation for pseudo-finite fields

- Let  $K$  be a pseudo-finite field.
- Want to understand definable sets  $X \subset K^n$ .
- Method: Associate **invariants** to  $X$ :
  - Dimension  $\dim(X)$
  - Measure  $\mu(X)$

# Situation for pseudo-finite fields

- Let  $K$  be a pseudo-finite field.
- Want to understand definable sets  $X \subset K^n$ .
- Method: Associate **invariants** to  $X$ :
  - Dimension  $\dim(X)$
  - Measure  $\mu(X)$

# Situation for pseudo-finite fields

- Let  $K$  be a pseudo-finite field.
- Want to understand definable sets  $X \subset K^n$ .
- Method: Associate **invariants** to  $X$ :
  - Dimension  $\dim(X)$
  - Measure  $\mu(X)$

# Dimension

## Definition (Dimension)

View  $X \subset K^n$  as a subset of  $\tilde{K}^n$ . Set  $\dim(X) :=$  dimension of Zariski-closure of  $X$  in  $\tilde{K}^n$ .

Example:  $X$  absolute irreducible variety

- $K$  is PAC  $\Rightarrow X(K)$  is dense in  $X(\tilde{K})$ .
- $\Rightarrow \dim X =$  alg. dimension of  $X$

# Dimension

## Definition (Dimension)

View  $X \subset K^n$  as a subset of  $\tilde{K}^n$ . Set  $\dim(X) :=$  dimension of Zariski-closure of  $X$  in  $\tilde{K}^n$ .

Example:  $X$  absolute irreducible variety

- $K$  is PAC  $\Rightarrow X(K)$  is dense in  $X(\tilde{K})$ .
- $\Rightarrow \dim X =$  alg. dimension of  $X$

# Dimension

## Definition (Dimension)

View  $X \subset K^n$  as a subset of  $\tilde{K}^n$ . Set  $\dim(X) :=$  dimension of Zariski-closure of  $X$  in  $\tilde{K}^n$ .

Example:  $X$  absolute irreducible variety

- $K$  is PAC  $\Rightarrow X(K)$  is dense in  $X(\tilde{K})$ .
- $\Rightarrow \dim X =$  alg. dimension of  $X$

# Dimension

## Definition (Dimension)

View  $X \subset K^n$  as a subset of  $\tilde{K}^n$ . Set  $\dim(X) :=$  dimension of Zariski-closure of  $X$  in  $\tilde{K}^n$ .

Example:  $X$  absolute irreducible variety

- $K$  is PAC  $\Rightarrow X(K)$  is dense in  $X(\tilde{K})$ .
- $\Rightarrow \dim X =$  alg. dimension of  $X$



# Dimension

## Definition (Dimension)

View  $X \subset K^n$  as a subset of  $\tilde{K}^n$ . Set  $\dim(X) :=$  dimension of Zariski-closure of  $X$  in  $\tilde{K}^n$ .

Example:  $X$  absolute irreducible variety

- $K$  is PAC  $\Rightarrow X(K)$  is dense in  $X(\tilde{K})$ .
- $\Rightarrow \dim X =$  alg. dimension of  $X$

# Dimension

## Definition (Dimension)

View  $X \subset K^n$  as a subset of  $\tilde{K}^n$ . Set  $\dim(X) :=$  dimension of Zariski-closure of  $X$  in  $\tilde{K}^n$ .

Example:  $X$  absolute irreducible variety

- $K$  is PAC  $\Rightarrow X(K)$  is dense in  $X(\tilde{K})$ .
- $\Rightarrow \dim X =$  alg. dimension of  $X$

# Measure: example

Example:  $X := \{\text{squares}\} := \{x^2 \mid x \in K\}$ .

- $\dim X = 1$
- “half of the elements” of  $K$  are squares.
- $\rightsquigarrow \mu(X) = \frac{1}{2}$ .

# Measure: example

Example:  $X := \{\text{squares}\} := \{x^2 \mid x \in K\}$ .

- $\dim X = 1$
- “half of the elements” of  $K$  are squares.
- $\rightsquigarrow \mu(X) = \frac{1}{2}$ .

# Measure: example

Example:  $X := \{\text{squares}\} := \{x^2 \mid x \in K\}$ .

- $\dim X = 1$
- “half of the elements” of  $K$  are squares.
- $\rightsquigarrow \mu(X) = \frac{1}{2}$ .

# Measure: example

Example:  $X := \{\text{squares}\} := \{x^2 \mid x \in K\}$ .

- $\dim X = 1$
- “half of the elements” of  $K$  are squares.
- $\rightsquigarrow \mu(X) = \frac{1}{2}$ .

# Measure: example

Example:  $X := \{\text{squares}\} := \{x^2 \mid x \in K\}$ .

- $\dim X = 1$
- “half of the elements” of  $K$  are squares.
- $\rightsquigarrow \mu(X) = \frac{1}{2}$ .

# Almost-quantifier-elimination

For definition of  $\mu$  need **almost-quantifier-elimination**:

## Theorem

*$K$  PSF field,  $X$  definable.  $\Rightarrow X = \pi(V(K))$ , where*

- *$V$  is algebraic*
- *$\pi$  is a projection*
- *$\pi \upharpoonright_V$  has finite fibers.*



# Almost-quantifier-elimination

For definition of  $\mu$  need **almost-quantifier-elimination**:

## Theorem

$K$  PSF field,  $X$  definable.  $\Rightarrow X = \pi(V(K))$ , where

- $V$  is algebraic
- $\pi$  is a projection
- $\pi \upharpoonright_V$  has finite fibers.

# Almost-quantifier-elimination

For definition of  $\mu$  need **almost-quantifier-elimination**:

## Theorem

$K$  PSF field,  $X$  definable.  $\Rightarrow X = \pi(V(K))$ , where

- $V$  is algebraic
- $\pi$  is a projection
- $\pi \upharpoonright_V$  has finite fibers.

# Almost-quantifier-elimination

For definition of  $\mu$  need **almost-quantifier-elimination**:

## Theorem

$K$  PSF field,  $X$  definable.  $\Rightarrow X = \pi(V(K))$ , where

- $V$  is algebraic
- $\pi$  is a projection
- $\pi \upharpoonright_V$  has finite fibers.

# Almost-quantifier-elimination

For definition of  $\mu$  need **almost-quantifier-elimination**:

## Theorem

$K$  PSF field,  $X$  definable.  $\Rightarrow X = \pi(V(K))$ , where

- $V$  is algebraic
- $\pi$  is a projection
- $\pi \upharpoonright_V$  has finite fibers.

# Almost-quantifier-elimination

For definition of  $\mu$  need **almost-quantifier-elimination**:

## Theorem

$K$  PSF field,  $X$  definable.  $\Rightarrow X = \pi(V(K))$ , where

- $V$  is algebraic
- $\pi$  is a projection
- $\pi \upharpoonright_V$  has finite fibers.

# Definition of measure

## Definition of $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

## Example: $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .



# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Definition of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .



# Defintion of measure

Definition of  $\mu(X)$ :

- $X \subset K^n$  algebraic set  $\rightsquigarrow \mu(X) :=$  number of irreducible components of Zariski closure of  $X$  in  $\tilde{K}$
- $X$  not algebraic:
  - Almost-ge  $\rightsquigarrow \pi: V \rightarrow X$
  - If all fibers have size  $n \rightsquigarrow \mu(X) := \frac{1}{n}\mu(V)$ .
  - Otherwise: A bit more work. (Best way: use Galois stratifications.)

Example:  $X = \{\text{squares}\}$ :

- $\{\text{squares}\} = \pi(V(y^2 = x))$  where  $\pi = \text{proj. on } x\text{-coordinate}$
- $V := V(y^2 = x)$  is absolutely irreducible  $\rightarrow \mu(V) = 1$ .
- Fiber have 2 elements except at  $x = 0$ .
- $\rightsquigarrow$  Indeed  $\mu(\{\text{squares}\}) = \frac{1}{2}$ .

# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Theorem (Hrushovski)

*These properties axiomatize the measure.*

# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Theorem (Hrushovski)

*These properties axiomatize the measure.*

# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Theorem (Hrushovski)

*These properties axiomatize the measure.*

# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Theorem (Hrushovski)

*These properties axiomatize the measure.*

# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Theorem (Hrushovski)

*These properties axiomatize the measure.*

# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Theorem (Hrushovski)

*These properties axiomatize the measure.*

# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Theorem (Hrushovski)

*These properties axiomatize the measure.*



# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Theorem (Hrushovski)

*These properties axiomatize the measure.*

# Properties

What does  $\mu$  satisfy?

- $\mu(V) = 1$  if  $V$  absolutely irreducible.
- $\mu(X \dot{\cup} Y) = \begin{cases} \mu(X) + \mu(Y) & \text{if } \dim X = \dim Y \\ \mu(X) & \text{if } \dim X > \dim Y \end{cases}$
- $\mu(X \times Y) = \mu(X) \cdot \mu(Y)$
- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- More generally: **Fubini**:  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

## Theorem (Hrushovski)

*These properties axiomatize the measure.*

# Outline

- 1 Measures on pseudo-finite fields
- 2 How to generalize?
- 3 Results

# Generalize: To which fields?

Generalize the measure to other fields. Which fields?

- Need control over definable sets.
- Recall:  $K$  PSF  $\iff K$  PAC, perfect and  $\text{Gal}_K := \text{Gal}(\tilde{K}/K) = \hat{\mathbb{Z}}$ .

Theorem (Hrushovski)

*Almost-qs still holds if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded.*

- $\text{Gal}_K$  **bounded** := has only a finite number of quotients of each fixed cardinality

Question (Hrushovski)

Does a measure exist if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded?

# Generalize: To which fields?

Generalize the measure to other fields. Which fields?

- Need control over definable sets.
- Recall:  $K$  PSF  $\iff K$  PAC, perfect and  $\text{Gal}_K := \text{Gal}(\tilde{K}/K) = \hat{\mathbb{Z}}$ .

Theorem (Hrushovski)

*Almost-qs still holds if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded.*

- $\text{Gal}_K$  **bounded** := has only a finite number of quotients of each fixed cardinality

Question (Hrushovski)

Does a measure exist if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded?

# Generalize: To which fields?

Generalize the measure to other fields. Which fields?

- Need control over definable sets.
- Recall:  $K$  PSF  $\iff K$  PAC, perfect and  $\text{Gal}_K := \text{Gal}(\tilde{K}/K) = \hat{\mathbb{Z}}$ .

Theorem (Hrushovski)

*Almost-qs still holds if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded.*

- $\text{Gal}_K$  **bounded** := has only a finite number of quotients of each fixed cardinality

Question (Hrushovski)

Does a measure exist if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded?

# Generalize: To which fields?

Generalize the measure to other fields. Which fields?

- Need control over definable sets.
- Recall:  $K$  PSF  $\iff K$  PAC, perfect and  $\text{Gal}_K := \text{Gal}(\tilde{K}/K) = \hat{\mathbb{Z}}$ .

Theorem (Hrushovski)

*Almost-ge still holds if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded.*

- $\text{Gal}_K$  **bounded** := has only a finite number of quotients of each fixed cardinality

Question (Hrushovski)

Does a measure exist if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded?

# Generalize: To which fields?

Generalize the measure to other fields. Which fields?

- Need control over definable sets.
- Recall:  $K$  PSF  $\iff K$  PAC, perfect and  $\text{Gal}_K := \text{Gal}(\tilde{K}/K) = \hat{\mathbb{Z}}$ .

## Theorem (Hrushovski)

*Almost- $q_e$  still holds if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded.*

- $\text{Gal}_K$  **bounded** := has only a finite number of quotients of each fixed cardinality

## Question (Hrushovski)

Does a measure exist if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded?



# Generalize: To which fields?

Generalize the measure to other fields. Which fields?

- Need control over definable sets.
- Recall:  $K$  PSF  $\iff K$  PAC, perfect and  $\text{Gal}_K := \text{Gal}(\tilde{K}/K) = \hat{\mathbb{Z}}$ .

## Theorem (Hrushovski)

*Almost-qs still holds if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded.*

- $\text{Gal}_K$  **bounded** := has only a finite number of quotients of each fixed cardinality

## Question (Hrushovski)

Does a measure exist if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded?

# Generalize: To which fields?

Generalize the measure to other fields. Which fields?

- Need control over definable sets.
- Recall:  $K$  PSF  $\iff K$  PAC, perfect and  $\text{Gal}_K := \text{Gal}(\tilde{K}/K) = \hat{\mathbb{Z}}$ .

## Theorem (Hrushovski)

*Almost-qs still holds if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded.*

- $\text{Gal}_K$  **bounded** := has only a finite number of quotients of each fixed cardinality

## Question (Hrushovski)

Does a measure exist if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded?

# Generalize: To which fields?

Generalize the measure to other fields. Which fields?

- Need control over definable sets.
- Recall:  $K$  PSF  $\iff K$  PAC, perfect and  $\text{Gal}_K := \text{Gal}(\tilde{K}/K) = \hat{\mathbb{Z}}$ .

## Theorem (Hrushovski)

*Almost-qs still holds if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded.*

- $\text{Gal}_K$  **bounded** := has only a finite number of quotients of each fixed cardinality

## Question (Hrushovski)

Does a measure exist if  $K$  PAC, perfect and  $\text{Gal}_K$  bounded?

# Generalized axioms

Can we expect all the axioms?

- **Fubini:**  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Example:  $K = \tilde{K}$ ,  $f: K^\times \rightarrow K^\times, x \mapsto x^2$ .

Fibers have 2 elements, but left and right set are the same. ⚡

# Generalized axioms

Can we expect all the axioms?

- **Fubini:**  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Example:  $K = \tilde{K}$ ,  $f: K^\times \rightarrow K^\times, x \mapsto x^2$ .

Fibers have 2 elements, but left and right set are the same. ⚡

# Generalized axioms

Can we expect all the axioms?

- **Fubini:**  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Example:  $K = \tilde{K}$ ,  $f: K^\times \rightarrow K^\times, x \mapsto x^2$ .

Fibers have 2 elements, but left and right set are the same. ⚡

# Generalized axioms

Can we expect all the axioms?

- **Fubini:**  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Example:  $K = \tilde{K}$ ,  $f: K^\times \rightarrow K^\times, x \mapsto x^2$ .

Fibers have 2 elements, but left and right set are the same. ⚡

# Generalized axioms

Can we expect all the axioms?

- **Fubini:**  $f: X \rightarrow Y$  definable, surjective, with constant fiber size  $\Rightarrow \mu(X) = \mu(Y) \cdot \mu(\text{fiber})$ .

Example:  $K = \tilde{K}$ ,  $f: K^\times \rightarrow K^\times, x \mapsto x^2$ .

Fibers have 2 elements, but left and right set are the same. ⚡



# Generalized axioms

Can we expect all the axioms?

- **Fubini**: No.

# Generalized axioms

Can we expect all the axioms?

- **Fubini:** No.

Instead:

- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- $\mu(X \times Y) = \mu(X) \times \mu(Y)$
- (• And as before  $\mu(\text{abs. irr}) = 1$  and  $\mu(X \dot{\cup} Y) = \dots$ )

These are our new axioms.

# Generalized axioms

Can we expect all the axioms?

- **Fubini:** No.

Instead:

- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- $\mu(X \times Y) = \mu(X) \times \mu(Y)$

(• And as before  $\mu(\text{abs. irr}) = 1$  and  $\mu(X \dot{\cup} Y) = \dots$ )

These are our new axioms.

# Generalized axioms

Can we expect all the axioms?

- **Fubini:** No.

Instead:

- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- $\mu(X \times Y) = \mu(X) \times \mu(Y)$
- (• And as before  $\mu(\text{abs. irr}) = 1$  and  $\mu(X \dot{\cup} Y) = \dots$ )

These are our new axioms.

# Generalized axioms

Can we expect all the axioms?

- **Fubini:** No.

Instead:

- $f: X \rightarrow Y$  definable bijection  $\Rightarrow \mu(X) = \mu(Y)$
- $\mu(X \times Y) = \mu(X) \times \mu(Y)$
- (• And as before  $\mu(\text{abs. irr}) = 1$  and  $\mu(X \dot{\cup} Y) = \dots$ )

These are our new axioms.

# Example: alg. closed fields

This works for  $K = \tilde{K}$ :

- $X$  definable  $\Rightarrow X$  constructible
- $\mu(X) :=$  number of irreducible components of  $X$  of maximal dimension.
- This satisfies all axioms.

# Example: alg. closed fields

This works for  $K = \tilde{K}$ :

- $X$  definable  $\Rightarrow X$  constructible
- $\mu(X) :=$  number of irreducible components of  $X$  of maximal dimension.
- This satisfies all axioms.

# Example: alg. closed fields

This works for  $K = \tilde{K}$ :

- $X$  definable  $\Rightarrow X$  constructible
- $\mu(X) :=$  number of irreducible components of  $X$  of maximal dimension.
- This satisfies all axioms.



# Example: alg. closed fields

This works for  $K = \tilde{K}$ :

- $X$  definable  $\Rightarrow X$  constructible
- $\mu(X) :=$  number of irreducible components of  $X$  of maximal dimension.
- This satisfies all axioms.

# Outline

- ① Measures on pseudo-finite fields
- ② How to generalize?
- ③ Results**

# Result

Let  $K$  be PAC, perfect and  $\text{Gal}_K$  bounded.

## Theorem (H.)

- 1 *If  $\text{Gal}_K$  is pro-cyclic, it works ( $\mu$  exists).*
- 2 *With one (natural) additional assumption, it is even unique.*
- 3 *For general bounded  $\text{Gal}_K$  (e.g.  $\text{Gal}_K = \hat{\mathbb{Z}} * \hat{\mathbb{Z}}$ ), no measure exists.*

# Result

Let  $K$  be PAC, perfect and  $\text{Gal}_K$  bounded.

## Theorem (H.)

- 1 *If  $\text{Gal}_K$  is pro-cyclic, it works ( $\mu$  exists).*
- 2 *With one (natural) additional assumption, it is even unique.*
- 3 *For general bounded  $\text{Gal}_K$  (e.g.  $\text{Gal}_K = \hat{\mathbb{Z}} * \hat{\mathbb{Z}}$ ), no measure exists.*

# Result

Let  $K$  be PAC, perfect and  $\text{Gal}_K$  bounded.

## Theorem (H.)

- 1 *If  $\text{Gal}_K$  is pro-cyclic, it works ( $\mu$  exists).*
- 2 *With one (natural) additional assumption, it is even unique.*
- 3 *For general bounded  $\text{Gal}_K$  (e.g.  $\text{Gal}_K = \hat{\mathbb{Z}} * \hat{\mathbb{Z}}$ ), no measure exists.*

# Result

Let  $K$  be PAC, perfect and  $\text{Gal}_K$  bounded.

## Theorem (H.)

- 1 *If  $\text{Gal}_K$  is pro-cyclic, it works ( $\mu$  exists).*
- 2 *With one (natural) additional assumption, it is even unique.*
- 3 *For general bounded  $\text{Gal}_K$  (e.g.  $\text{Gal}_K = \hat{\mathbb{Z}} * \hat{\mathbb{Z}}$ ), no measure exists.*

# Why only for pro-cyclic Galois groups

- Does not work in general because there are many definable bijections:

Have more or less  $X \xrightarrow{1:1} Y$  for  $Y \subsetneq X$ .

- Does work for pro-cyclic Galois group, because:
  - It works for PSF fields and...
  - language of  $K$  (with  $\text{Gal}_K$  pro-cyclic) is a “simplification” of language of PSF.

# Why only for pro-cyclic Galois groups

- Does not work in general because there are many definable bijections:

Have more or less  $X \xrightarrow{1:1} Y$  for  $Y \subsetneq X$ .

- Does work for pro-cyclic Galois group, because:
  - It works for PSF fields and...
  - language of  $K$  (with  $\text{Gal}_K$  pro-cyclic) is a “simplification” of language of PSF.



# Why only for pro-cyclic Galois groups

- Does not work in general because there are many definable bijections:  
Have more or less  $X \xrightarrow{1:1} Y$  for  $Y \subsetneq X$ .
- Does work for pro-cyclic Galois group, because:
  - It works for PSF fields and...
  - language of  $K$  (with  $\text{Gal}_K$  pro-cyclic) is a “simplification” of language of PSF.

# Why only for pro-cyclic Galois groups

- Does not work in general because there are many definable bijections:  
Have more or less  $X \xrightarrow{1:1} Y$  for  $Y \subsetneq X$ .
- Does work for pro-cyclic Galois group, because:
  - It works for PSF fields and...
  - language of  $K$  (with  $\text{Gal}_K$  pro-cyclic) is a “simplification” of language of PSF.

# Why only for pro-cyclic Galois groups

- Does not work in general because there are many definable bijections:  
Have more or less  $X \xrightarrow{1:1} Y$  for  $Y \subsetneq X$ .
- Does work for pro-cyclic Galois group, because:
  - It works for PSF fields and...
  - language of  $K$  (with  $\text{Gal}_K$  pro-cyclic) is a “simplification” of language of PSF.

# Why only for pro-cyclic Galois groups

- Does not work in general because there are many definable bijections:  
Have more or less  $X \xrightarrow{1:1} Y$  for  $Y \subsetneq X$ .
- Does work for pro-cyclic Galois group, because:
  - It works for PSF fields and...
  - language of  $K$  (with  $\text{Gal}_K$  pro-cyclic) is a “simplification” of language of PSF.

# Why only for pro-cyclic Galois groups

- Does not work in general because there are many definable bijections:  
Have more or less  $X \xrightarrow{1:1} Y$  for  $Y \subsetneq X$ .
- Does work for pro-cyclic Galois group, because:
  - It works for PSF fields and...
  - language of  $K$  (with  $\text{Gal}_K$  pro-cyclic) is a “simplification” of language of PSF.

# When $\text{Gal}_K$ pro-cyclic

Example: Consider  $X = \{\text{squares}\}$ .

- If  $K$  is PSF, then  $K = \{\text{squares}\} \dot{\cup} \{\text{non-squares}\}$ .  
 $\mu_K(\{\text{squares}\}) = \mu_K(\{\text{non-squares}\}) = \frac{1}{2}$ .
- Now add all square roots to  $K$ :  $L := K \cup \sqrt{K} \cup \sqrt{\sqrt{K}} \cup \dots$
- $\rightsquigarrow$  All non-squares also become squares.
- $\rightsquigarrow \mu_L(\{\text{squares}\}) = \mu_K(\{\text{squares}\}) + \mu_K(\{\text{non-squares}\}) = 1$

# When $\text{Gal}_K$ pro-cyclic

Example: Consider  $X = \{\text{squares}\}$ .

- If  $K$  is PSF, then  $K = \{\text{squares}\} \dot{\cup} \{\text{non-squares}\}$ .  
 $\mu_K(\{\text{squares}\}) = \mu_K(\{\text{non-squares}\}) = \frac{1}{2}$ .
- Now add all square roots to  $K$ :  $L := K \cup \sqrt{K} \cup \sqrt{\sqrt{K}} \cup \dots$
- $\rightsquigarrow$  All non-squares also become squares.
- $\rightsquigarrow \mu_L(\{\text{squares}\}) = \mu_K(\{\text{squares}\}) + \mu_K(\{\text{non-squares}\}) = 1$

# When $\text{Gal}_K$ pro-cyclic

Example: Consider  $X = \{\text{squares}\}$ .

- If  $K$  is PSF, then  $K = \{\text{squares}\} \dot{\cup} \{\text{non-squares}\}$ .  
 $\mu_K(\{\text{squares}\}) = \mu_K(\{\text{non-squares}\}) = \frac{1}{2}$ .
- Now add all square roots to  $K$ :  $L := K \cup \sqrt{K} \cup \sqrt{\sqrt{K}} \cup \dots$
- $\rightsquigarrow$  All non-squares also become squares.
- $\rightsquigarrow \mu_L(\{\text{squares}\}) = \mu_K(\{\text{squares}\}) + \mu_K(\{\text{non-squares}\}) = 1$



# When $\text{Gal}_K$ pro-cyclic

Example: Consider  $X = \{\text{squares}\}$ .

- If  $K$  is PSF, then  $K = \{\text{squares}\} \dot{\cup} \{\text{non-squares}\}$ .  
 $\mu_K(\{\text{squares}\}) = \mu_K(\{\text{non-squares}\}) = \frac{1}{2}$ .
- Now add all square roots to  $K$ :  $L := K \cup \sqrt{K} \cup \sqrt{\sqrt{K}} \cup \dots$
- $\rightsquigarrow$  All non-squares also become squares.
- $\rightsquigarrow \mu_L(\{\text{squares}\}) = \mu_K(\{\text{squares}\}) + \mu_K(\{\text{non-squares}\}) = 1$

# When $\text{Gal}_K$ pro-cyclic

Example: Consider  $X = \{\text{squares}\}$ .

- If  $K$  is PSF, then  $K = \{\text{squares}\} \dot{\cup} \{\text{non-squares}\}$ .  
 $\mu_K(\{\text{squares}\}) = \mu_K(\{\text{non-squares}\}) = \frac{1}{2}$ .
- Now add all square roots to  $K$ :  $L := K \cup \sqrt{K} \cup \sqrt{\sqrt{K}} \cup \dots$
- $\rightsquigarrow$  All non-squares also become squares.
- $\rightsquigarrow \mu_L(\{\text{squares}\}) = \mu_K(\{\text{squares}\}) + \mu_K(\{\text{non-squares}\}) = 1$

# When $\text{Gal}_K$ pro-cyclic

Example: Consider  $X = \{\text{squares}\}$ .

- If  $K$  is PSF, then  $K = \{\text{squares}\} \dot{\cup} \{\text{non-squares}\}$ .  
 $\mu_K(\{\text{squares}\}) = \mu_K(\{\text{non-squares}\}) = \frac{1}{2}$ .
- Now add all square roots to  $K$ :  $L := K \cup \sqrt{K} \cup \sqrt{\sqrt{K}} \cup \dots$
- $\rightsquigarrow$  All non-squares also become squares.
- $\rightsquigarrow \mu_L(\{\text{squares}\}) = \mu_K(\{\text{squares}\}) + \mu_K(\{\text{non-squares}\}) = 1$

# 1. When $\text{Gal}_K$ pro-cyclic

General idea is the same.

- Pretend that  $K$  is an extension of a PSF field.
- Which definable sets in the PSF field become part of  $X$  when extending to  $K$ ?
- Set  $\mu(X)$  to be the sum of the measures of those sets.
- Real way to do this is using Galois stratification.

Remark: If one tries to define  $\mu$  for  $K$  directly (without going through the PSF-measure), it does not work.

# 1. When $\text{Gal}_K$ pro-cyclic

General idea is the same.

- Pretend that  $K$  is an extension of a PSF field.
- Which definable sets in the PSF field become part of  $X$  when extending to  $K$ ?
- Set  $\mu(X)$  to be the sum of the measures of those sets.
- Real way to do this is using Galois stratification.

Remark: If one tries to define  $\mu$  for  $K$  directly (without going through the PSF-measure), it does not work.

# 1. When $\text{Gal}_K$ pro-cyclic

General idea is the same.

- Pretend that  $K$  is an extension of a PSF field.
- Which definable sets in the PSF field become part of  $X$  when extending to  $K$ ?
  - Set  $\mu(X)$  to be the sum of the measures of those sets.
  - Real way to do this is using Galois stratification.

Remark: If one tries to define  $\mu$  for  $K$  directly (without going through the PSF-measure), it does not work.

# 1. When $\text{Gal}_K$ pro-cyclic

General idea is the same.

- Pretend that  $K$  is an extension of a PSF field.
- Which definable sets in the PSF field become part of  $X$  when extending to  $K$ ?
- Set  $\mu(X)$  to be the sum of the measures of those sets.
- Real way to do this is using Galois stratification.

Remark: If one tries to define  $\mu$  for  $K$  directly (without going through the PSF-measure), it does not work.

# 1. When $\text{Gal}_K$ pro-cyclic

General idea is the same.

- Pretend that  $K$  is an extension of a PSF field.
- Which definable sets in the PSF field become part of  $X$  when extending to  $K$ ?
- Set  $\mu(X)$  to be the sum of the measures of those sets.
- Real way to do this is using Galois stratification.

Remark: If one tries to define  $\mu$  for  $K$  directly (without going through the PSF-measure), it does not work.



# Uniqueness without Fubini

Example:  $K$  is PSF and  $X = \{\text{squares}\}$ .

- We have a definable bijection  $f: \{\text{squares}\} \rightarrow \{\text{non-squares}\}$ :
  - Take  $a \in \{\text{non-squares}\}$ . Then  $f(x) := a \cdot x$ .

So  $\mu(\{\text{squares}\}) = \mu(\{\text{non-squares}\})$ .

- $\mu(\{\text{squares}\}) + \mu(\{\text{non-squares}\}) = \mu(K) = 1$
- $\Rightarrow \mu(\{\text{squares}\}) = \frac{1}{2}$ .

The general idea is the same.

# Uniqueness without Fubini

Example:  $K$  is PSF and  $X = \{\text{squares}\}$ .

- We have a definable bijection  $f: \{\text{squares}\} \rightarrow \{\text{non-squares}\}$ :
  - Take  $a \in \{\text{non-squares}\}$ . Then  $f(x) := a \cdot x$ .

So  $\mu(\{\text{squares}\}) = \mu(\{\text{non-squares}\})$ .

- $\mu(\{\text{squares}\}) + \mu(\{\text{non-squares}\}) = \mu(K) = 1$
- $\Rightarrow \mu(\{\text{squares}\}) = \frac{1}{2}$ .

The general idea is the same.

# Uniqueness without Fubini

Example:  $K$  is PSF and  $X = \{\text{squares}\}$ .

- We have a definable bijection  $f: \{\text{squares}\} \rightarrow \{\text{non-squares}\}$ :

- Take  $a \in \{\text{non-squares}\}$ . Then  $f(x) := a \cdot x$ .

So  $\mu(\{\text{squares}\}) = \mu(\{\text{non-squares}\})$ .

- $\mu(\{\text{squares}\}) + \mu(\{\text{non-squares}\}) = \mu(K) = 1$
- $\Rightarrow \mu(\{\text{squares}\}) = \frac{1}{2}$ .

The general idea is the same.

# Uniqueness without Fubini

Example:  $K$  is PSF and  $X = \{\text{squares}\}$ .

- We have a definable bijection  $f: \{\text{squares}\} \rightarrow \{\text{non-squares}\}$ :
  - Take  $a \in \{\text{non-squares}\}$ . Then  $f(x) := a \cdot x$ .

So  $\mu(\{\text{squares}\}) = \mu(\{\text{non-squares}\})$ .

- $\mu(\{\text{squares}\}) + \mu(\{\text{non-squares}\}) = \mu(K) = 1$
- $\Rightarrow \mu(\{\text{squares}\}) = \frac{1}{2}$ .

The general idea is the same.

# Uniqueness without Fubini

Example:  $K$  is PSF and  $X = \{\text{squares}\}$ .

- We have a definable bijection  $f: \{\text{squares}\} \rightarrow \{\text{non-squares}\}$ :
  - Take  $a \in \{\text{non-squares}\}$ . Then  $f(x) := a \cdot x$ .

So  $\mu(\{\text{squares}\}) = \mu(\{\text{non-squares}\})$ .

- $\mu(\{\text{squares}\}) + \mu(\{\text{non-squares}\}) = \mu(K) = 1$
- $\Rightarrow \mu(\{\text{squares}\}) = \frac{1}{2}$ .

The general idea is the same.

# Uniqueness without Fubini

Example:  $K$  is PSF and  $X = \{\text{squares}\}$ .

- We have a definable bijection  $f: \{\text{squares}\} \rightarrow \{\text{non-squares}\}$ :

- Take  $a \in \{\text{non-squares}\}$ . Then  $f(x) := a \cdot x$ .

So  $\mu(\{\text{squares}\}) = \mu(\{\text{non-squares}\})$ .

- $\mu(\{\text{squares}\}) + \mu(\{\text{non-squares}\}) = \mu(K) = 1$
- $\Rightarrow \mu(\{\text{squares}\}) = \frac{1}{2}$ .

The general idea is the same.

# Uniqueness without Fubini

Example:  $K$  is PSF and  $X = \{\text{squares}\}$ .

- We have a definable bijection  $f: \{\text{squares}\} \rightarrow \{\text{non-squares}\}$ :
  - Take  $a \in \{\text{non-squares}\}$ . Then  $f(x) := a \cdot x$ .

So  $\mu(\{\text{squares}\}) = \mu(\{\text{non-squares}\})$ .

- $\mu(\{\text{squares}\}) + \mu(\{\text{non-squares}\}) = \mu(K) = 1$
- $\Rightarrow \mu(\{\text{squares}\}) = \frac{1}{2}$ .

The general idea is the same.

# Uniqueness without Fubini

Example:  $K$  is PSF and  $X = \{\text{squares}\}$ .

- We have a definable bijection  $f: \{\text{squares}\} \rightarrow \{\text{non-squares}\}$ :
  - Take  $a \in \{\text{non-squares}\}$ . Then  $f(x) := a \cdot x$ .

So  $\mu(\{\text{squares}\}) = \mu(\{\text{non-squares}\})$ .

- $\mu(\{\text{squares}\}) + \mu(\{\text{non-squares}\}) = \mu(K) = 1$
- $\Rightarrow \mu(\{\text{squares}\}) = \frac{1}{2}$ .

The general idea is the same.